



ALLIANCE SEED AAC LEROY VB



AAC LeRoy VB has the amazing combination of high yields, disease resistance, consistent harvest quality **AND** midge tolerance that makes it a go-to variety for farmers across Western Canada.

AAC LeRoy VB strengths:

- Higher yielding than all of the checks, yielding 113% of Carberry and 110% of Unity.
- Maturity 2 days earlier than Carberry, 1 day later than Unity
- Lodging score better than Unity, and within range of checks
- Height 6 cm shorter than Unity
- Kernel weight higher than all of the checks
- Excellent disease resistance
- Resistance to Wheat Midge

AAC LeRoy VB neutral traits:

- Test weight within range of the checks

AAC LeRoy VB weaknesses:

- Grain protein concentration 0.6 units lower than Carberry
- MS reaction to leaf spot

Parentage: BW431/Carberry



CWRS

2025 MCVET CWRS Wheat Trials

Variety	Yield bu/ac	Protein %	Maturity +/- 99 days	Height +/- 81 cm	Spike Awned	Lodging	Sprouting	Loose Smut	Common Bunt	Leaf Spots 1	Stem Rust	Leaf Rust	Stripe Rust	FHB 2
AAC Brandon (check)	74	14.4	2	0	Y	VG	P	MR	S	I	R	R	MR	MR
AAC Stoughton VB	76	13.2	3	2	Y	G	-	-	MR	-	R	R	I	MR
AAC Wheatland VB	77	14.4	1	0	Y	VG	G	R	MR	S	R	R	I	I
AAC Westking	76	14.3	2	-3	Y	VG	F	-	R	-	MR	R	I	MR
AAC Starbuck VB	77	14.7	1	1	Y	G	F	MR	S	S	I	MR	MR	MR
AAC Hodge VB	77	14.1	2	8	Y	VG	P	MS	R	R	R	R	R	MR
AAC Elie	72	14.5	2	-3	Y	VG	F	I	I	I	R	R	MR	I
AAC LeRoy VB	76	14.4	0	7	Y	G	G	-	I	MS	MR	MR	MR	MR

¹ Leaf Spots rating is a combination of resistance to tan spot, spot blotch and septoria leaf disease complex. ² Fusarium head blight (FHB) infection is highly influenced by genetics, environment and heading date. Under high levels of disease pressure, all varieties will experience yield and/or quality loss.

M=Medium, L=Late, VL=Very late, G=Good, VG = Very good, P=Poor, VP= Very Poor, F=Fair, R=Resistant, MR=Moderately Resistant,

I=Intermediate Resistance, MS= Moderately Susceptible, S=Susceptible

2025 SKVPG CWRS Wheat Trials

Variety	Yield (%)		Protein (%)	Resistance to									Stem Solid- ness	Rel. Maturity (days)	Seed wt (mg)	Volume Wt. (kg/hL)	Ht (cm)
	Area 1 & 2	Area 3 & 4		Lodging	Sprouting	Stem Rust	Leaf Rust	Stripe Rust	Loose Smut	Bunt	Leaf Spot	FHB					
AAC Brandon (check)	100	100	14.3	F	P	R	R	MR	MR	S	I	MR	H	101	35.7	80.7	81
AAC Hodge VB	102	107	-0.3	F	P	R	R	R	-	R	-	MR	H	-1	-1.4	0.3	6
AAC Wheatland VB	104	106	-0.2	G	G	R	R	I	R	MR	S	I	H	0	-0.7	0.1	1
AAC Starbuck VB	104	108	-0.2	F	F	I	MR	MR	MR	S	S	MR	H	0	-0.2	0.4	2
AAC Elie	99	99	0	F	F	R	R	MR	I	I	I	I	H	1	-0.7	-0.1	-2
AAC LeRoy VB	98	102	-0.1	F	G	MR	MR	MR	-	I	MS	MR	H	-1	-0.5	0.3	6

M=Medium, L=Late, VL=Very late, G=Good, VG = Very good, P=Poor, VP= Very Poor, F=Fair, R=Resistant, MR=Moderately Resistant,

I=Intermediate Resistance, MS= Moderately Susceptible, S=Susceptible

2025 ABRVT CWRS Wheat Trials

Variety	Overall Yield	Low < 77 (bu/ac)	High ≥ 77 (bu/ac)	Maturity Rating (days +/-)	Protein (%)	Test Wt. (lb/bu)	TKW (g)	Height (cm)	Awn (y/n)	Lodging	Sprouting	Bunt	Stripe Rust	FHB
AAC Brandon (check)	100	100	100	104	14.0	63	39	84	Y	F	P	S	MR	MR
AAC Wheatland VB	104	104	104	0	-0.5	63	40	86	Y	VG	VG	MR	I	I
Stettler	97	98	97	0	0.1	63	38	92	Y	F	G	MR	MR	MS
AAC Hodge VB	103	101	105	-1	-0.3	63	37	91	Y	F	P	R	R	MR
AAC Starbuck VB	103	104	102	0	-0.2	63	39	87	Y	G	F	S	MR	MR
AAC LeRoy VB	100	101	99	0	-0.2	63	39	88	Y	G	G	I	MR	MR

M=Medium, L=Late, VL=Very late, G=Good, VG = Very good, P=Poor, VP= Very Poor, F=Fair, R=Resistant, MR=Moderately Resistant,

I=Intermediate Resistance, MS= Moderately Susceptible, S=Susceptible