

AAC Redberry is legendary for premium harvest quality, its beautiful red berries **AND** its early maturity. If that's what you are looking for, AAC Redberry remains one of the best options!

AAC Redberry strengths:

- High grain yield with earlier maturity
- Low DON accumulation
- Semi-dwarf stature
- High test weight
- R to: Stem rust, Leaf rust, Stripe rust and Loose smut.

AAC Redberry neutral traits:

 Good standability, less than AAC Viewfield or AAC Connery

AAC Redberry weakness:

MS to leaf spots

Parentage: Stettler/Glenn



2025 MCVET CWRS Wheat Trials

Variety	Yield bu/ac	Protein %	Maturity +/- 99 days	Height +/- 81 cm	Spike Awned	Lodging	Sprouting	Loose Smut	Common Bunt	Leaf Spots 1	Stem Rust	Leaf Rust	Stripe Rust	FHB 2
AAC Brandon (check)	74	14.4	2	0	Υ	VG	Р	MR	S		R	R	MR	MR
AAC Viewfield	75	14.5	3	-5	Υ	VG	G	S	MR	_	R	MR	R	- 1
AAC Hockley	73	14.4	2	-1	Υ	VG	F	-	R	-	MR	R	R	MR
AAC Wheatland VB	77	14.4	1	0	Υ	VG	G	R	MR	S	R	R	- 1	- 1
AAC Starbuck VB	77	14.7	1	1	Υ	G	F	MR	S	S	- 1	MR	MR	MR
CDC Envy	73	14.3	-1	2	Υ	G	F	-	R		- 1	R	MR	1
AAC Redberry	73	14.5	0	8	Υ	G	G	R	_	MS	R	R	R	_

1 Leaf Spots rating is a combination of resistance to tan spot, spot blotch and septoria leaf disease complex. 2 Fusarium head blight (FHB) infection is highly influenced by genetics, environment and heading date. Under high levels of disease pressure, all varieties will experience yield and/or quality loss.

M=Medium, L=Late, VL=Very late, G=Good, VG = Very good, P=Poor, VP= Very Poor, F=Fair, R=Resistant, MR=Moderately Resistant,

l=Intermediate Resistance, MS= Moderately Susceptible, S=Susceptible

2025 SKVPG CWRS Wheat Trials

	Yie	ld (%)	Protein	Resistance to									Stem	Rel.	Seed	Volume	
Variety	Area 1 & 2	Area 3 & 4	(%)	Lodging	Sprouting	Stem Rust	Leaf Rust	Stripe Rust	Loose Smut	Bunt	Leaf Spot	FHB	Solid- ness	Maturity (days)	wt (mg)	Wt. (kg/hL)	Ht (cm)
AAC Brandon (check)	100	100	14.3	F	Р	R	R	MR	MR	S	1	MR	Н	101	35.7	80.7	81
AAC Viewfield	104	101	-0.3	G	G	R	MR	R	S	MR	- 1	-1	Н	0	-2.2	0.8	-3
AAC Hodge VB	102	107	-0.3	F	Р	R	R	R		R		MR	Н	-1	-1.4	0.3	6
AAC Hockley	100	103	0.1	G	F	MR	R	R		R		MR	Н	0	-1.8	0.9	0
AAC Wheatland VB	104	106	-0.2	G	G	R	R	1	R	MR	S	-1	Н	0	-0.7	0.1	1
AAC Starbuck VB	104	108	-0.2	F	F	_	MR	MR	MR	S	S	MR	Н	0	-0.2	0.4	2
CDC Envy	98	100	-0.3	F	F	_	R	MR		R		- 1	Н	-2	0.5	-1.8	2
AAC Redberry	99	100	0	F	G	R	R	R	R	1	MS	1	н	-2	-1.7	0.7	6

M=Medium, L=Late, VL=Very late, G=Good, VG = Very good, P=Poor, VP= Very Poor, F=Fair, R=Resistant, MR=Moderately Resistant,

I=Intermediate Resistance, MS= Moderately Susceptible, S=Susceptible

2025 ABRVT CWRS Wheat Trials

Variety	Overall Yield	Low < 77 (bu/ac)	High ≥ 77 (bu/ac)	Maturity Rating (days +/-)	Protein (%)	Test Wt. (lb/bu)	TKW (g)	Height (cm)	Awn (y/n)	Lodging	Sprouting	Bunt	Stripe Rust	FHB
AAC Brandon (check)	100	100	100	104	14.0	63	39	84	Y	F	Р	S	MR	MR
AAC Viewfield	102	98	105	0	-0.3	63	37	80	Y	VG	G	MR	R	- 1
AAC Hockley	99	96	102	1	0.1	64	35	82	Y	VG	F	R	R	MR
AAC Wheatland VB	104	104	104	0	-0.5	63	40	86	Y	VG	VG	MR	- 1	1
Stettler	97	98	97	0	0.1	63	38	92	Y	F	G	MR	MR	MS
AAC Hodge VB	103	101	105	-1	-0.3	63	37	91	Y	F	Р	R	R	MR
AAC Starbuck VB	103	104	102	0	-0.2	63	39	87	Y	G	F	S	MR	MR
AAC Westking	102	99	104	0	-0.1	63	41	82	Y	VG	VG	R	_	MR
CDC Envy	96	91	101	-2	-0.4	61	38	85	Y	Р	G	R	MR	- 1
AAC Redberry	94	94	94	-3	-0.3	63	41	90	Υ	F	VG	1	R	1

M=Medium, L=Late, VL=Very late, G=Good, VG=Very good, P=Poor, VP=Very Poor, F=Fair, R=Resistant, MR=Moderately Resistant, I=Intermediate Resistance, MS= Moderately Susceptible, S=Susceptible